

ATOMIC-RESOLUTION STRUCTURE OF CYANOBACTRIAL CYTOCHROME C₆ WITH AN UNUSUAL SEQUENCE INSERTION

S. Krzywda^{1*}, **W. Bialek**², **A. Szczepaniak**², and **M. Jaskólski**¹

¹ Faculty of Chemistry, Adam Mickiewicz University, Poznań, Poland

² Faculty of Biotechnology, University of Wrocław, Poland

Keywords: high-resolution structure, cytochrome, photosynthesis

*) e-mail: szymon@amu.edu.pl

During photosynthesis, electron transfer between two membrane-bound complexes, cytochrome *b₆f* and photosystem I can be accomplished by the copper-containing protein, plastocyanin, or the heme protein, cytochrome *c₆* [1]. Cytochromes *c₆* are water-soluble, low-spin heme-containing proteins involved in the high-potential (340–390 mV) electron transport chain. They are characterized by low molecular mass, 80–90 amino acid residues in the mature protein, and have a covalently bound heme group.

Cytochrome *c₆* from the mesophilic cyanobacterium *Synechococcus* sp. PCC 7002 is unique among all known cytochromes *c₆* due to the presence of an unusual seven-residue insertion, K₄₄DGSKSL₅₀. Furthermore, the present protein is unusual because of its very high content (36%) of the smallest residues (glycine and alanine) [2].

The crystallization experiments used the hanging-drop vapor-diffusion method. Red crystals (Fig. 1) suitable for X-ray analysis were obtained from 10 mM sodium Hepes, pH 6.2, and 2.2 M ammonium sulfate over a period of one week.



Figure 1. Crystals of reduced cytochrome *c₆* from *Synechococcus* sp. PCC 7002.

The structure of the reduced form of cytochrome *c₆* has been determined at 1.2 Å and refined to an *R* factor of 0.107. It reveals that the overall fold of the protein is similar to that of other class I *c*-type cytochromes despite

the presence of the specific insertion. The insertion is located within the most variable region of cytochrome *c₆* sequence, *i.e.* between helices II and III. The first six residues (K₄₄DGSKS₄₉) form a loop, whereas the last residue, Leu₅₀, extends the N-terminal beginning of helix III. Several specific non-covalent interactions are found inside the insertion as well as between the insertion and the rest of the protein. Energetically significant cation... π interactions have been detected between Tyr₄₃, Tyr₅₆, Phe₆₈ and, respectively, Lys₃₇, Lys₄₈ and Arg₇₁ (Fig. 2).




Figure 2. N-H... π interaction between the only Arg and Phe68 ($2F_o - F_c$ electron density map contoured at 1.2 σ).

The crystal structure contains three copies of the *c₆* molecule per asymmetric unit and is characterized by unusually high packing density, with solvent occupying barely 17.58% of the crystal volume.

Acknowledgements: Some of the calculations were carried out in the Poznan Metropolitan Supercomputing and Networking Center. This work was funded in part by grants number N N204 245635 and N N303 3856 33 from the Ministry of Science and Higher Education.

References

- [1] W.A. Cramer, P.N. Furbacher, A. Szczepaniak, G.S. Tae, *Electron transport between photosystem II and photosystem I* (Academic Press, San Diego 1991).
- [2] W. Bialek, M. Nelson, K. Tamiola, T. Kallas, A. Szczepaniak, "Deeply branching *c₆*-like cytochromes of cyanobacteria", *Biochemistry* **47** (2008) 5515–5522.